

Dissecting d -Cubes into Smaller d -Cubes

Matthew Hudelson

*Department of Pure and Applied Mathematics, Washington State University,
Pullman, Washington 99164-3113*

Communicated by the Managing Editors

Received March 18, 1997

In this paper, we explore the following question: Given integers d and k , is it possible to subdivide a d -dimensional cube into k smaller d -dimensional cubes? In particular, we investigate bounds on the integer $c(d)$ which is the smallest integer

View metadata, citation and similar papers at core.ac.uk

given k , the asymptotic behavior of $c(d)$ for those d such that $\gcd(2^d - 1, k^d - 1) = 1$. Specifically, we show that if $\gcd(2^d - 1, 3^d - 1)$ then $c(d) < 6^d$ and that if $\gcd(2^d - 1, k^d - 1)$ then $c(d) = O((2k)^d)$. Finally, we derive the general asymptotic bound $c(d) = O((2d)^{d-1})$ which improves the currently known bound of $c(d) = O((2d)^d)$. © 1998 Academic Press, Inc.

1. BACKGROUND

Croft, Falconer, and Guy [1] discuss the following problem which has been attributed to Hadwiger and proposed by Fine and Niven [3]: *Given d , for which integers k can the unit d -dimensional cube Q_d be decomposed into k smaller d -cubes?* If we let D_d represent the set of all such integers, then let $c(d)$ be the smallest integer such that if $k \geq c(d)$ then $k \in D_d$.

In order to investigate $c(d)$ further, we prove the following elementary result.

THEOREM 1. *Suppose A is a set of positive integers $\{a_1, \dots, a_{2^d-1}\}$ such that, for each k , Q^d can be subdivided into a_k d -cubes and such that for each $m \in \{0, \dots, 2^d - 2\}$ there is a k_m such that $a_{k_m} \equiv m \pmod{2^d - 1}$. Then $c(d) \leq \max(A) - 2^d + 2$.*

By definition, we cannot subdivide Q_d into $c(d) - 1$ cubes. Suppose $c(d) > \max(A) - 2^d + 2$. There exists some $a \in A$ such that $a \equiv c(d) - 1 \pmod{2^d - 1}$, so we can write $c(d) - 1 = p(2^d - 1) + a$ for some integer p . If $p = 0$, then we can subdivide Q_d into $a = c(d) - 1$ cubes which is a contradiction. If $p > 0$, then we subdivide Q_d into a cubes, then subdivide one

of the smaller cubes into 2^d still smaller cubes, and then repeat this latter step $p-1$ more times to yield a subdivision of Q_d into $c(d)-1$ cubes which is again a contradiction. Therefore, $p < 0$. It follows that

$$c(d) - 1 = p(2^d - 1) + a \leq p(2^d - 1) + \max(A) \leq \max(A) - (2^d - 1)$$

and so $c(d) \leq \max(A) - 2^d + 2$, which contradicts our original assumption. The result follows. ■

2. BOUNDS ON $c(d)$ FOR $d \leq 5$

2.1. The Cases when $d \leq 3$

Since it is possible to subdivide a line segment into any number of smaller line segments, $c(1) = 1$, and in fact, $D_1 = \mathbf{N}$. In Fig. 1, we demonstrate that $c(2) \leq 6$ by showing how to subdivide a square into one, six, and eight smaller squares, respectively. In this case, we have $A = \{6, 7, 8\}$ although $A = \{1, 6, 8\}$ or $A = \{4, 6, 8\}$ would also serve to show that $c(2) \leq 6$. It is not possible to subdivide a square into two, three, or five smaller squares, so, in fact, $c(2) = 6$ and $D_2 = \mathbf{N} \setminus \{2, 3, 5\}$.

Exact values of $c(d)$ are unknown for $d > 2$, although it is conjectured that $c(3) = 48$. To see that $c(3) \leq 48$, we show that $A = \{1, 20, 38, 39, 49, 51, 54\}$ is a subset of D_3 . Then Theorem 1 gives us $c(3) \leq 54 - 8 + 2 = 48$. Figures 2 and 3 show how to subdivide a cube into a smaller cubes for each $a \in A$.

The subdivision of a cube into 54 smaller cubes is the most intricate and was discovered by Rychener and Zbinden (see [1]).

2.2. The Four-Dimensional Case

We show that $c(4) \leq 809$ by showing

$$\{1, 66, 131, 370, 435, 500, 634, 672, 693, 699, 737, 758, 764, 802, 823\} \subset D_4.$$

To show this, we first note that subdividing a cube into 3^4 cubes and then coalescing 2^4 of these into a cube results in a subdivision of a cube




Fig. 1. Subdividing a square into smaller squares, $a \in \{1, 6, 8\}$.




Fig. 2. Subdividing a cube into smaller cubes, $a \in \{1, 20, 38, 49\}$.

into 66 smaller cubes. Moreover, this strategy can be used to show that if $k \in D_4$, then $k + 65 \in D_4$. With this in mind, it suffices to show that $\{1, 370, 634, 672, 693\} \subset D_4$. Clearly, $a_1 = 1 \in D_4$ as we can leave the cube in one piece. Also, it is easy to see that $a_0 = 370 = 5^4 - 4^4 + 1 \in D_4$ and $a_2 = 672 = 6^4 - 5^4 + 1 \in D_4$.

Next, we note that $10^4 = 14 \cdot 5^4 + 2 \cdot 4^4 + 8 \cdot 2^4 + 610 \cdot 1^4$, and if we can assemble these cubes into one cube, the result is a subdivision of Q_4 into $a_4 = 634$ smaller cubes. Table I contains a list of “least corners” (a, b, c, d) for the “ k^4 ” hypercubes, $k \geq 2$, in the subdivision. The other corners of each hypercube are $(a + a', b + b', c + c', d + d')$, a', b', c' , and d' each in $\{0, k\}$.

Finally, to show $a_3 = 693 \in D_4$, we note that $10^4 = 13 \cdot 5^4 + 5 \cdot 3^4 + 53 \cdot 2^4 + 622 \cdot 1^4$. The corresponding list of least corners is contained in Table II.

2.3. An Upper Bound for $c(5)$

At this point, we demonstrate that $c(5) \leq 1891$, a bound credited to William Scott by Croft *et al.* [1, page 85]. We reproduce this bound to introduce a strategy which we will generalize in Section 3.

Our basic strategy will be, given $i \in \{0, 1, \dots, 30\}$, to subdivide Q_5 into 32 cubes with half the edge length ($1/2$ -cubes) and then to subdivide i of




Fig. 3. Subdividing a cube into smaller cubes, $a \in \{39, 51, 54\}$.

these each into $3^5 = 243$ $1/6$ -cubes. As a result, we will have subdivided Q_d into $(3^5 - 1)i + (2^5)$ smaller cubes. Since $\gcd(2^5 - 1, 3^5 - 1) = 1$, it follows that distinct choices i yield distinct choices of j such that $(3^5 - 1)i \equiv j \pmod{2^5 - 1}$. Furthermore, if we make a judicious choice of which i of the $1/2$ -cubes to further subdivide, we will be able to recombine a number of groups of 2^5 of the $1/6$ -cubes into $1/3$ -cubes.

TABLE I
Least Corners for the $a_4 = 634$ Case

"5 ⁴ " hypercubes:	(0, 0, 0, 0)	(0, 0, 0, 5)	(0, 0, 5, 0)	(0, 0, 5, 5)
	(0, 5, 0, 0)	(0, 5, 0, 5)	(0, 5, 5, 0)	(0, 5, 5, 5)
	(5, 0, 0, 0)	(5, 0, 0, 5)	(5, 0, 5, 0)	(5, 0, 5, 5)
	(5, 5, 0, 0)	(5, 5, 0, 5)		
"4 ⁴ " hypercubes:	(5, 5, 5, 0)	(5, 5, 5, 4)		
"2 ⁴ " hypercubes:	(5, 5, 5, 8)	(5, 5, 7, 8)	(5, 7, 5, 8)	(5, 7, 7, 8)
	(7, 5, 5, 8)	(7, 5, 7, 8)	(7, 7, 5, 8)	(7, 7, 7, 8)

TABLE II
Least Corners for the $a_3 = 693$ Case

“5 ⁴ ” hypercubes:	(0, 0, 0, 0)	(0, 0, 0, 5)	(0, 0, 5, 0)	(0, 0, 5, 5)
	(0, 5, 0, 0)	(0, 5, 0, 5)	(0, 5, 5, 0)	(0, 5, 5, 5)
	(5, 0, 0, 0)	(5, 0, 0, 5)	(5, 0, 5, 0)	(5, 0, 5, 5)
	(5, 5, 0, 0)			
“3 ⁴ ” hypercubes:	(5, 5, 7, 0)	(5, 5, 7, 3)	(5, 5, 7, 6)	(5, 5, 0, 7)
	(5, 5, 3, 7)			
“2 ⁴ ” hypercubes:	(5, 5, 0, 5)	(5, 7, 0, 5)	(7, 5, 0, 5)	(7, 7, 0, 5)
	(5, 5, 2, 5)	(5, 7, 2, 5)	(7, 5, 2, 5)	(7, 7, 2, 5)
	(5, 5, 4, 5)	(5, 7, 4, 5)	(7, 5, 4, 5)	(7, 7, 4, 5)
	(5, 5, 5, 0)	(5, 7, 5, 0)	(7, 5, 5, 0)	(7, 7, 5, 0)
	(5, 5, 5, 2)	(5, 7, 5, 2)	(7, 5, 5, 2)	(7, 7, 5, 2)
	(8, 5, 0, 7)	(8, 7, 0, 7)	(5, 8, 0, 7)	(8, 5, 2, 7)
	(8, 7, 2, 7)	(5, 8, 2, 7)	(8, 5, 4, 7)	(8, 7, 4, 7)
	(5, 8, 4, 7)	(8, 5, 8, 0)	(8, 7, 8, 0)	(5, 8, 8, 0)
	(8, 5, 8, 2)	(8, 7, 8, 2)	(5, 8, 8, 2)	(8, 5, 6, 4)
	(8, 7, 6, 4)	(5, 8, 6, 4)	(8, 5, 8, 4)	(8, 7, 8, 4)
	(5, 8, 8, 4)	(8, 5, 6, 6)	(8, 7, 6, 6)	(5, 8, 6, 6)
	(8, 5, 8, 6)	(8, 7, 8, 6)	(5, 8, 8, 6)	(8, 5, 6, 8)
	(8, 7, 6, 8)	(5, 8, 6, 8)	(8, 5, 8, 8)	(8, 7, 8, 8)
	(5, 8, 8, 8)			

Suppose $i = 2^k$. Then we subdivide a $2^k \times 1^{5-k}$ slab of 1/2-cubes into 1/6-cubes and recombine $6^k 2^{5-k}$ of these into 1/3-cubes. Therefore, we have subdivided Q_d into $(2^5 - 2^k)$ 1/2-cubes, 3^k 1/3-cubes, and $6^k(3^{5-k} - 2^{5-k})$ 1/6-cubes.

If $i = \sum_p 2^{k_p}$, where each k_p is a distinct nonnegative integer, then for each p , we subdivide a $2^{k_p} \times 1^{5-k_p}$ slab of 1/2-cubes into 1/6 cubes and recombine each slab as in the previous paragraph. But we can also recombine across slab boundaries to further reduce the total number of cubes into which Q_d is subdivided. In Section 3, we will generalize this recombination argument. There, we introduce the function

$$\psi\left(\sum_p 2^{k_p}\right) = \sum_p 2^p 3^{k_p}$$

where $k_0 > k_1 > \dots > k_m \geq 0$. By using the more general results in Section 3, we see that the total number of each type of cube is $(2^5 - i)$ 1/2-cubes, $\psi(i)$ 1/3-cubes, and $(3^5 i - 2^5 \psi(i))$ 1/6-cubes. Thus, the total number of cubes is $\tau(i) = (3^5 - 1)i - (2^5 - 1)\psi(i) + 2^5$. Table III contains the values of $\psi(i)$ and $\tau(i)$ for $0 \leq i \leq 30$.

The largest value for $\tau(i)$ in Table III is 1921, so by Theorem 1, $c(5) \leq 1921 - 32 + 2 = 1891$.

TABLE III

Values of $\psi(i)$ and $\tau(i)$ for $0 \leq i \leq 30$

i	$\psi(i)$	$\tau(i)$	i	$\psi(i)$	$\tau(i)$	i	$\psi(i)$	$\tau(i)$
0	0	32	11	37	1547	22	111	1915
1	1	243	12	45	1541	23	119	1909
2	3	423	13	49	1659	24	135	1655
3	5	603	14	57	1653	25	139	1773
4	9	721	15	65	1647	26	147	1767
5	11	901	16	81	1393	27	155	1761
6	15	1019	17	83	1573	28	171	1507
7	19	1137	18	87	1691	29	179	1501
8	27	1131	19	91	1809	30	195	1247
9	29	1311	20	99	1803			
10	33	1429	21	103	1921			

3. ASYMPTOTIC UPPER BOUNDS FOR $c(d)$

3.1. The Case when $\gcd(2^d - 1, 3^d - 1) = 1$

We generalize the construction for $d=5$ to the general case when $\gcd(2^d - 1, 3^d - 1) = 1$. For such a d , we define each slab S_k , $k \in \{0, 1, \dots, d-1\}$, by

$$S_k = [0, 6)^k \times [0, 3) \times [3, 6)^{d-k-1}.$$

We make our cubes and slabs half open on the higher side to make arguments concerning disjoint unions easier to formulate. We note that S_k is then the disjoint union of 2^k half-open cubes, each three units on a side. We also note that if $j < k$, then $S_j \cap S_k$ is empty since the $k+1$ term of S_j is $[3, 6)$ while the $k+1$ term of S_k is $[0, 3)$.

Next, we define

$$R_k = [0, 6)^k \times [0, 2) \times [4, 6)^{d-k-1}.$$

As an observation, $R_k \subset S_k$ for each k . Next, given a decreasing set of non-negative integers $X = \{k_0, k_1, \dots, k_q\}$, we define N_X to be the cartesian product expansion of R_{k_q} with the $(k_0 + 1)$, $(k_1 + 1)$, and so on to the $(k_{q-1} + 1)$ terms replaced with $[0, 4)$. If $X = \{k_0\}$, then $N_X = R_{k_0}$. Here, we note that N_X is the disjoint union of $3^k 2^q$ half-open cubes, each two units on a side.

Finally, we define, for the same set X as before, $X_i = \{k_0, \dots, k_i\}$, and

$$C_X = \bigsqcup_{i=0}^p N_{X_i},$$

where \bigsqcup denotes a disjoint union. We have a disjoint union since if $i > j$, then the $(k_i + 1)$ term of N_{X_i} is $[4, 6)$, while the $(k_i + 1)$ term of N_{X_j} is $[0, 4)$.

We now claim that $C_X \subset \bigsqcup k \in X S_k$. If $x \in C_X$, then $x = (x_1, \dots, x_d) \in N_{X_m}$ for some m . Then

$$x_i \in \begin{cases} [0, 6) & \text{if } i \leq k_m \\ [0, 2) & \text{if } i = k_m + 1 \\ [0, 4) & \text{if } i \in \{k_0 + 1, k_1 + 1, \dots, k_{m-1} + 1\} \\ [4, 6) & \text{otherwise.} \end{cases}$$

If there is no $k_i \in X_{m-1}$ such that $x_{k_i+1} \in [0, 3)$, then $x \in S_{k_m}$ and we are done. Otherwise, let k_q be the smallest element of X_{m-1} such that $x_{k_q+1} \in [0, 3)$. We then have that

$$x_i \in \begin{cases} [0, 6) & \text{if } i \leq k_q \\ [0, 3) & \text{if } i = k_q + 1 \\ [3, 6) & \text{otherwise,} \end{cases}$$

and so $x \in S_{m_q}$ and we are done with the claim.

Now, if $\gcd(2^d - 1, 3^d - 1) = 1$, for each $m \in \{0, 1, \dots, 2^d - 2\}$, $(3^d - 1)m$ is distinct modulo $2^d - 1$. Therefore, we can subdivide Q_d into 2^d smaller cubes, and subdivide m of these smaller cubes into 3^d still smaller cubes to arrive at a subdivision of Q_d into $2^d + m(3^d - 1)$ cubes.

We improve this simple dissection by using slabs and recoalescing into larger cubes. Suppose $m = \sum_{i=0}^p 2^{k_i}$ with the k_i 's decreasing. Then we dissect $6[0, 1)^d$ into 2^d three-unit cubes, but we recoalesce the appropriate three-unit cubes into slabs $S_{k_0}, S_{k_1}, \dots, S_{k_p}$, leaving $2^d - m$ three-unit cubes. We then dissect each of these slabs into 3^d one-unit cubes.

Above, we showed that, for $X_i = \{k_0, \dots, k_i\}$ and $X = X_p$, $C_X \subset \bigsqcup_{k \in X} S_k$ and that $C_X = \bigsqcup_i N_{X_i}$. Furthermore, we saw that N_{X_i} is the disjoint union of $3^{k_i} 2^i$ two-unit cubes. Therefore, C_X is the disjoint union of $\sum_{i=0}^p 3^{k_i} 2^i$ two-unit cubes. We call this sum $\psi(m)$ to reflect its dependance on m .

Then, we can recoalesce $2^d \psi(m)$ one-unit cubes of $\bigsqcup_{k \in X} S_k$ into two-unit cubes. As a result, we have dissected $6Q_d$ into $(2^d - m)$ three-unit cubes, $\psi(m)$ two-unit cubes, and $(m3^d - \psi(m)2^d)$ one-unit cubes for a total of

$2^d + (3^d - 1)m - (2^d - 1)\psi(m)$ cubes in the entire dissection. This dissection has $(2^d - 1)\psi(m)$ fewer cubes than the simple dissection. For $m \in \{0, 1, \dots, 2^d - 2\}$, $2^d + (3^d - 1)m - (2^d - 1)\psi(m) < 6^d$. Also, for each $i \in \{0, 1, \dots, 2^d - 2\}$, there is some m such that $2^d + (3^d - 1)m - (2^d - 1)\psi(m) \equiv 1 \pmod{2^d - 1}$ since $\gcd(2^d - 1, 3^d - 1) = 1$. Therefore, from Theorem 1, we may conclude that $c(d) < 6^d$. Computer experiments show that for $d \leq 25$ that in the case $\gcd(2^d - 1, 3^d - 1) = 1$ that $c(d) < 6^d/4.1$.

3.2. General Asymptotics

THEOREM 2. *If $\gcd(2^d - 1, k^d - 1) = 1$ for infinitely many values of d , then $c(d) = O((2k)^{d-1})$.*

If k is even, then $\gcd(2^d - 1, k^d - 1) = \gcd(2^d - 1, (k/2)^d - 1)$, so we assume k is odd. We generate each element a_i in A by first subdividing Q_d into 2^d cubes and then subdividing i of these cubes into k^d smaller cubes. At this point, we have subdivided Q_d into $2^d + i(k^d - 1)$ cubes. Suppose $i = \sum_{p=1}^m 2^{n_p}$. Then we choose our i cubes in $2^{n_p} \times 1^{d-n_p}$ slabs. For each slab, we then recombine groups of 2^d small cubes into cubes having twice the edge length. In the " n_p " slab, we may recombine $(2k)^{n_p} (k - 1)^{d-n_p}$ small cubes in this manner, leaving $(2k)^{n_p} (k^{d-n_p} - (k - 1)^{d-n_p})$ small cubes unmolested. Then if we compute the number of small cubes in all of Q_d , we obtain

$$\begin{aligned}
 \# \text{ of small cubes} &= \sum_{p=1}^m (2k)^{n_p} (k^{d-n_p} - (k-1)^{d-n_p}) \\
 &\leq \sum_{n=0}^{d-1} (2k)^n (k^{d-n} - (k-1)^{d-n}) \\
 &= \sum_{n=0}^{d-1} (2k)^n \left(\sum_{q=0}^{d-n-1} k^q (k-1)^{d-n-1-q} \right) \\
 &\leq \sum_{n=0}^{d-1} (2k)^n (d-n-1) k^{d-n-1} \\
 &\leq \sum_{n=0}^{d-1} (2k)^{d-1} (d-n-1) 2^{-(d-n-1)} \\
 &\leq (2k)^{d-1} \sum_{n=0}^{d-1} n 2^{-n} \\
 &\leq (2k)^{d-1} \sum_{n=0}^{\infty} n 2^{-n} \\
 &= 4(2k)^{d-1}
 \end{aligned}$$

which is an $O((2k)^{d-1})$ quantity. The number of recoalesced cubes is bounded by k^d and the number of cubes into which Q_d was originally decomposed is bounded by 2^d . It follows that for each i , we can choose $a_i \in A$ such that

$$a_i \leq \left(4 + \frac{k}{2^{d-1}} + \frac{2}{k^{d-1}}\right) (2k)^{d-1}.$$

This last quantity is $O((2k)^{d-1})$ and independent of i . This implies $c(d) = O((2k)^{d-1})$ as desired. ■

We now prove our general asymptotic bound for all d , namely that $c(d) = O((2d)^{d-1})$. In order to do this we need two lemmas, the first of which appears in Erdős [2] and the second of which is an application of elementary calculus.

LEMMA 1. *For any positive integer d , $\gcd(2^d - 1, 3^d - 1, \dots, (d+1)^d - 1) = 1$.*

The following proof appears in Erdős [2]. Suppose this statement is not true. Then there is some prime number p which is a factor of $k^d - 1$ for each $k \in A = \{1, 2, 3, \dots, d+1\}$. We then note that $k^d \equiv 1 \pmod p$ for such integers k . Further, we note that $p > (d+1)$, since if not then p is a factor of $p^d - 1$ which is absurd. Therefore, A is a set of $d+1$ distinct elements of the group \mathbf{Z}_p^+ under multiplication, all of whose orders divide d . Since this group is cyclic, there can be at most d such elements which is a contradiction. ■

LEMMA 2. *There is some constant C such that $S(m) = \sum_{k=1}^{m-1} (1 - k/m)^{m-2} < C$ for all integers $m > 2$.*

The k th term of $S(m)$ has limit e^{-k} as m increases without bound, so each term has some constant c_k which bounds it. What we do not know is whether we can find a set of such constants $\{c_1, c_2, \dots\}$ such that the series $\sum c_k$ converges. We show that, for $k \geq 4$, e^{-k} in fact bounds $(1 - k/m)^{m-2}$ from above, and so we may choose

$$\begin{aligned} C &= c_1 + c_2 + c_3 + \sum_{k=4}^{\infty} e^{-k} \\ &= c_1 + c_2 + c_3 + e^{-4}(1 - e^{-1})^{-1}. \end{aligned}$$

For $k \geq 4$, we consider the function $f(x) = (1 - k/x)^{x-2}$. We seek to show that $f(x)$ is increasing for $x \geq k+1$. This will suffice to show that, on $[k+1, \infty)$, $f(x)$ is bounded by $\lim_{x \rightarrow \infty} f(x) = e^{-k}$. We compute $f'(x) = (1 - k/x)^{x-3} ((1 - k/x) \ln(1 - k/x) + (x-2)k/x^2)$ which we wish to show is

positive for $x \geq k+1$. Since $1 - k/x > 0$ for $x \geq k+1$, it is enough to show that $g(x) = \ln(1 - k/x) + k(x-2)/(x^2(1 - k/x)) > 0$. Expanding each term of $g(x)$ in a power series of $1/x$, we obtain

$$\begin{aligned}\ln(1 - k/x) &= \sum_{i=1}^{\infty} -(k^i)/(ix^i) \\ kx/x^2(1 - k/x) &= \sum_{i=1}^{\infty} k^i/x^i \\ 2k/x^2(1 - k/x) &= \sum_{i=2}^{\infty} 2k^{i-1}/x^i\end{aligned}$$

and, since these series all converge absolutely when $x \geq k+1$, we have that

$$g(x) = \sum_{i=2}^{\infty} \frac{(i-1)k^i - 2ik^{i-1}}{ix^i}.$$

The numerator of each term is $k^{i-1}(k(i-1) - 2i)$. Since $k \geq 4$ and $i \geq 2$, this expression is at least $4^{i-1}(4i - 4 - 2i) = 4^{i-1}(2i - 4) \geq 0$ with equality only when $i = 2$. Therefore, $g(x)$ is positive, and then so is $f'(x)$. Hence, $f(x)$ is increasing for $x \in [k+1, \infty)$. This completes the proof of the lemma. ■

THEOREM 3. $c(d) = O((2d)^{d-1})$.

We begin by subdividing Q_d into 2^d mini-cubes. Since $2^d \geq d-1$ for all positive d , we may subdivide $d-1$ of the mini-cubes each into 2^d micro-cubes. Now, from Lemma 1, given an $a \in \{0, 1, \dots, 2^d - 2\}$, there is some $(d-1)$ -tuple $(a_3, a_4, \dots, a_{d+1})$ such that $0 \leq a_i \leq 2^d - 2$ and such that $1 + \sum_{k=3}^{d+1} a_k(k^d - 1) \equiv a \pmod{2^d - 1}$. With this in mind, for each k , we subdivide a_k of the micro-cubes in the $(k-2)$ nd mini-cube into k^d nano-cubes. At this point, the entire cube Q_d has been subdivided into $2^d - d + 1$ mini-cubes, $(d-1)2^d - \sum_{k=3}^{d+1} a_k$ micro-cubes, and $\sum_{k=3}^{d+1} a_k k^d$ nano-cubes. Summing these, we have that Q_d is subdivided into

$$1 + (d-1)(2^d - 1) + \sum_{k=3}^{d+1} a_k(k^d - 1)$$

smaller cubes. This last quantity is equivalent to a modulo $2^d - 1$.

Now, if the $a_k = \sum_{p=1}^m 2^{n_p}$ nano-cubes in the $(k-2)$ nd micro-cube are arranged in $2^{n_p} \times 1^{d-n_p}$ slabs, as in the proof of Theorem 2, then we may recombine many of these nano-cubes into cubes having twice the edge length. By the computation in Theorem 2, there will be at most $4(2k)^{d-1}$ nano-cubes in the $(k-2)$ nd micro-cube remaining after the recombining

process. Furthermore, the total number of cubes into which the $(k-2)$ nd micro-cube has been subdivided satisfies

$$\text{number of cubes} \leq \left(4 + \frac{d+1}{2^{d-1}} + \frac{2}{3^{d-1}}\right) (2k)^{d-1} \leq \frac{37}{6} (2k)^{d-1}.$$

This follows from the last formula in the proof of Theorem 2, noting that $3 \leq k \leq d+1$.

It follows that the total number of cubes into which Q_d has been subdivided satisfies

$$\begin{aligned} \#(\text{cubes}) &\leq 1 + (d-1)(2^d - 1) + \sum_{k=3}^{d+1} \frac{37}{6} (2k)^{d-1} \\ &\leq 1 + (d-1)(2^d - 1) + \frac{37}{6} (2(d+1))^{d-1} \sum_{k=3}^{d+1} \left(\frac{k}{d+1}\right)^{d-1} \\ &\leq (2d)^{d-1} \left((2d)^{1-d} + 2d^{2-d} + \frac{37}{6} eC \right), \end{aligned}$$

where C is the constant in Lemma 2 bounding $S(d+1)$ which is greater than $\sum_{k=3}^{d+1} (k/(d+1))^{d-1}$. This last quantity is $O((2d)^{d-1})$, which proves the result. ■

REFERENCES

1. H. T. Croft, K. J. Falconer, and R. K. Guy, "Unsolved Problems in Geometry," Springer-Verlag, New York, 1991.
2. P. Erdős, Remarks on some problems in number theory, in "Proc., Conference of Balkan Mathematics, Belgrade," June 1974.
3. N. J. Fine and I. Niven, Problem E724, *Amer. Math. Monthly* **53** (1946), 271; solution, **54** (1947) 41–42.
4. R. K. Guy, Monthly research problems, *Amer. Math. Monthly* **82** (1975), 1001.
5. R. K. Guy, Monthly research problems, *Amer. Math. Monthly* **84** (1977), 809–810.
6. C. Meier, Decomposition of a cube into smaller cubes, *Amer. Math. Monthly* **81** (1974), 630–632.